


Models for Estimation of Chemical Distribution and Fate in Response to Remedial Alternatives in the Lower Willamette River

Technical Briefing Portland Harbor Superfund Project

Bruce Hope, Ph.D.
Air Quality Division
Oregon Department of Environmental Quality
Portland, Oregon



Purpose

- * Evaluate impact of source control activities and remedial action alternatives on chemical levels in:
 - Sediment and water
 - Fish
- * Provide estimates (by segment) of:
 - Loads from external sources
 - Ambient levels in relation to PRGs
 - Probability of site recontamination
 - Contributions from outside the site (e.g., upstream)
 - Effects of monitored natural recovery
 - Downstream responses to actions upstream

Response to remedial actions

- * How much? Source reductions may not translate into a proportional reduction in sediment, water, fish
- * How fast? Temporal response of system will depend on which source is targeted for remediation
- * How permanent? If ultimate source is not remediated, potential for recontamination will remain
- * How extensive (spatially)? Downstream extent of response depends on both biological and physical processes
- * How necessary? Considering the role of “natural” attenuation in load reduction

A decision support tool

- * A computer model, in conjunction with data analyses, may be used to quantitatively and comparatively address these primary study questions
- * Use of a model may decrease uncertainty in effectiveness of the preferred remedy
- * Benefits
 - Constrain, synthesize and interpret data
 - Compare remedial alternatives on same basis
- * Drawbacks
 - Level of uncertainty in results may be unacceptable

Key specifications

- * TRANSPARENCY

- * Link transport & fate \Rightarrow food web models

- * Incorporate spatial variability

- Exposure of mobile species

- * Incorporate temporal variability

- Seasonal variation; Non-steady-state dynamics

- * Evaluate organics and other chemicals

- * Results as mass or concentration


- * Food web-specific:

- Invertebrates as whole animals, intra- and up-trophic consumption, model fish \Rightarrow bird \Rightarrow egg

Concept


- * Linked biotic and abiotic sub-models
 - After Gobas et al. (1998) - Fraser River
- * Abiotic - transport & fate sub-model
 - After Davis (2003, 2004) for San Francisco Bay
 - Considers back- and cross-flows between segments
 - STELLA®
- * Biotic - food web sub-model
 - After Arnot & Gobas (2004)
 - Same generic structure in every segment
 - Multi-segment exposure for mobile species
 - Excel® & Visual Basic®

Changes in water & sediment concentrations




Constant load (0.0001 kg d^{-1})
to water in segments 1,2,3

Changes in sediment concentration across segments




Constant load (0.0001 kg d⁻¹)
to water in segments 1,2,3

Changes in sediment concentration across segments




Constant load (0.0001 kg d⁻¹)
to water in segment 16

PCB-118 tissue residues over time (baseline)




Constant load (0.0001 kg d^{-1})
to water in segments 1,2,3

PCB-118 tissue residues over time (post-remedy)




Constant load to water
ceases at $t = 600$ d

Changes in water & sediment concentration (periodic)




Periodic load (0.01 kg d⁻¹ every 365 d)
to water in segments 1,2,3

PCB-118 tissue residues over time (periodic)




Periodic load (0.01 kg d^{-1} every 365 d)
to water in segments 1,2,3

Changes in water & sediment concentration (one load)




One load (0.1 kg d⁻¹ at day 600)
to water in segments 1,2,3

PCB-118 tissue residues over time (one load)




One load (0.1 kg d^{-1} at day 600)
to water in segments 1,2,3


River segments




Segments mapped to the physical river



Transport & fate sub-model for a single river segment




Abiotic sub-model relationships between segments



Sub-model linkages over multiple segments

FOOD WEB (BIOTIC) SUB-MODEL




TRANSPORT & FATE (ABIOTIC) SUB-MODEL

Technical Briefing (May 2006)


Food web sub-model for a single segment

Exposure to sediment [S], sediment porewater [P], overlying water [W]



Technical Briefing (May 2006)

Data flow and manipulation between sub-models



Contaminants

* POLYCYCLIC AROMATIC HYDROCARBONS

- Naphthalene
- Phenanthrene
- Fluoranthrene
- Benz(a)anthracene
- Benzo(b)fluoranthene
- Dibenzo(a,h)anthracene
- Benzo(ghi)perylene

* METALS

- Mercury (methylmercury)
- Arsenic
- Other

* POLYCHLORINATED BIPHENYLS

- PCB 18
- PCB 66
- **PCB 118**
- PCB 153
- PCB 194

* DIOXINS

- TCDD, PCDD, HCDD

* PESTICIDES

- DDT
- DDE
- DDD

Most sensitive parameters

- ~* Henry's Law constant
- ~* Octanol-water partition coefficient
 - * Water temperature
- ~* River flow rate
 - * Active sediment depth
 - * Concentration of solids in sediment
 - * Density of particles (suspended) in water
 - * Sediment solids burial rate
- ~* Bioavailable concentration in water (+ related terms)
 - * Water content fraction of organism (~ lipids)
 - * Dietary fraction(s)